



Drug Facts

Active ingredient (in each chewable tablet) **Purpose**
Bismuth subsalicylate 262 mg ... Upset stomach reliever and antidiarrheal

Uses relieves • travelers' diarrhea • diarrhea
• upset stomach due to overindulgence in food and drink, including:
• heartburn • indigestion • nausea • gas • belching • fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have • an ulcer • a bleeding problem
• bloody or black stool

Ask a doctor before use if you have • fever • mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for
• anticoagulation (thinning the blood) • diabetes • gout • arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if • symptoms get worse or last more than 2 days • ringing in the ears or loss of hearing occurs
• diarrhea lasts more than 2 days

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- fully chew before swallowing
- adults and children 12 years and over:
 - 2 chewable tablets every $\frac{1}{2}$ hour or 4 chewable tablets every hour as needed for diarrhea
 - 2 chewable tablets every $\frac{1}{2}$ hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
 - do not exceed 8 doses (16 chewable tablets) in 24 hours
 - use until diarrhea stops but not more than 2 days
 - children under 12 years: ask a doctor
 - drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information • each chewable tablet contains:

- calcium 50 mg • salicylate 111 mg • low sodium
- Store at no greater than 25°C. Avoid excessive heat (over 40°C.)

Inactive ingredients acacia, beeswax, calcium carbonate, caprylic/capric triglyceride, carmine, carnauba wax, citric acid, corn starch, corn syrup, dextrin, FD&C Blue No. 2 Aluminum Lake, flavor, hydrogenated coconut oil, maltodextrin, potassium hydroxide, propylene glycol, shellac, sodium benzoate, sorbic acid, sorbitol, soy lecithin, sucrose, titanium dioxide, water

Questions? 1-800-717-3786