


HEAVITAS REGULAR

Helvetica Neue Condensed Bold
 Helvetica Neue LT Std 87 Heavy Condensed
 Avenir Next Condensed Regular

HEX:1E75BA K: 0
 C: 84.77 Y: 1.17
 M: 49.61
 C: 47.07 M: 76.94
 C: 71.83 M: 85.28
 HEX:5F3F72 K: 12.62
 C: 47.07 M: 76.94
 C: 71.83 M: 85.28
 HEX:94589C K: 0
 C: 47.07 Y: 5.22
 M: 76.94
 C: 71.83 M: 85.28
 HEX:1E75BA K: 0
 C: 84.77 Y: 1.17
 M: 49.61

DIELINE KEY

4.4375"





3 735811 10224 2

THIS PRODUCT IS PACKAGED IN A CHILD-RESISTANT AND TAMPER EVIDENT PACKAGE. USE ONLY IF BLISTERS ARE INTACT.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Distributed by **Wellspring**
Affordable Trusted Relief

Albion, NY 10852 U.S.A.
 866-933-6337
 wellspringmeds.com


24 SOFTGELS**


✓ Suitable for Adults and Children 12+
 ✓ Controls Symptoms of Diarrhea


Loperamide HCl 2 mg

SYMPTOM CONTROL

ANTI-DIARRHEAL








24 SOFTGELS**

Made in USA*



ANTI-DIARRHEAL

Loperamide HCl 2 mg

Drug Facts	Drug Facts (continued)
<p>Active ingredient (in each softgel) Purpose</p> <p>Loperamide HCl 2 mg.....Anti-diarrheal</p>	<p>of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p>
<p>Use</p> <p>Controls symptoms of diarrhea, including Travelers' Diarrhea</p>	<p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.</p>
<p>Warnings</p> <p>Allergy alert: Do not use if you have ever had a rash or other allergic reaction to Loperamide HCl.</p> <p>Heart alert: Taking more than directed can cause serious heart problems or death.</p> <p>Do not use:</p> <ul style="list-style-type: none"> ▪ If you have bloody or black stool ▪ In children under 12 years of age <p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> ▪ Fever ▪ Mucus in the stool ▪ A history of liver disease ▪ A history of abnormal heart rhythm <p>Ask a doctor before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.</p> <p>When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ▪ Symptoms get worse ▪ Diarrhea lasts more than 2 days ▪ You get abdominal swelling or bulging. These may be signs ▶ 	<p>Directions</p> <ul style="list-style-type: none"> ▪ Drink plenty of clear fluids to help prevent dehydration caused by diarrhea ▪ Not for use in children under 12 years of age ▪ Adults and children 12 years and over: 2 softgels after the first loose stool; 1 softgel after each subsequent loose stool; but no more than 4 softgels in 24 hours <p>Other information</p> <ul style="list-style-type: none"> ▪ Do not use if blister unit is torn or open ▪ Store at 20° - 25°C (68° - 77°F) <p>Inactive ingredients</p> <p>butylated hydroxyanisole, FD&C Blue #1, gelatin, glycerin, glyceryl caprylate, polyoxyl 40 hydrogenated castor oil, purified water.</p> <p>Questions? call toll-free 1-866-933-6337</p>

*Made in USA with globally sourced ingredients
 **Each liquid-filled capsule contains 2 mg loperamide HCl

3"