

<p><b>Drug Facts</b></p> <p><b>Active ingredients (in each 30 mL) Purpose</b></p> <p>Acetaminophen 650 mg ..... Pain reliever/fever reducer</p> <p>Dextromethorphan HBr 30 mg ..... Cough suppressant</p> <p>Doxylamine succinate 12.5 mg ..... Antihistamine</p>	<p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> </ul>	<p><b>If pregnant or breast-feeding</b>, ask a health professional before use. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>						
<p><b>Uses</b> temporarily relieves common cold/flu symptoms:</p> <ul style="list-style-type: none"> <li>cough due to minor throat and bronchial irritation</li> <li>sore throat</li> <li>headache</li> <li>minor aches and pains</li> <li>fever</li> <li>runny nose and sneezing</li> </ul>	<p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>liver disease</li> <li>glaucoma</li> <li>cough that occurs with too much phlegm (mucus)</li> <li>a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema</li> <li>trouble urinating due to enlarged prostate gland</li> </ul>	<p><b>Directions</b></p> <ul style="list-style-type: none"> <li>take only as directed</li> <li>only use the dose cup provided</li> <li>do not exceed 4 doses per 24 hrs</li> </ul> <table border="1"> <tr> <td>adults &amp; children 12 yrs &amp; over</td> <td>30 mL per dose every 6 hrs</td> </tr> <tr> <td>children 4 to under 12 yrs</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 yrs</td> <td>do not use</td> </tr> </table>	adults & children 12 yrs & over	30 mL per dose every 6 hrs	children 4 to under 12 yrs	ask a doctor	children under 4 yrs	do not use
adults & children 12 yrs & over	30 mL per dose every 6 hrs							
children 4 to under 12 yrs	ask a doctor							
children under 4 yrs	do not use							
<p><b>Warnings</b></p> <p><b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> <li>more than 4 doses (30 mL each dose) in 24 hours, which is the maximum daily amount for this product</li> <li>with other drugs containing acetaminophen</li> <li>3 or more alcoholic drinks every day while using this product</li> </ul> <p><b>Allergy Alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> <li>skin reddening</li> <li>blisters</li> <li>rash</li> </ul> <p>If a skin reaction occurs, stop use and seek medical help right away</p> <p><b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p>	<p><b>Ask a doctor or pharmacist before use if you are</b></p> <ul style="list-style-type: none"> <li>taking sedatives or tranquilizers</li> <li>taking the blood thinning drug warfarin</li> </ul>	<p><b>Other information</b></p> <ul style="list-style-type: none"> <li>each 30 mL contains: sodium 28.23 mg</li> <li>store at no greater than 25°C and do not refrigerate</li> </ul>						
	<p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>excitability may occur, especially in children</li> <li>marked drowsiness may occur</li> <li>avoid alcoholic drinks</li> <li>be careful when driving a motor vehicle or operating machinery</li> <li>alcohol, sedatives, and tranquilizers may increase drowsiness</li> </ul>	<p><b>Inactive ingredients</b> citric acid, FD&amp;C blue no. 1, FD&amp;C red no. 40, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum</p>						
	<p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>pain or cough gets worse or lasts more than 7 days</li> <li>fever gets worse or lasts more than 3 days</li> <li>redness or swelling is present</li> <li>new symptoms occur</li> <li>cough comes back or occurs with rash or headache that lasts.</li> </ul> <p>These could be signs of a serious condition.</p>	<p><b>Questions? 1-800-362-1683</b></p>						



**FAST  
RELIEF**  
**HBP**

VICKS



**NyQuil**<sup>TM</sup>

**HIGH BLOOD  
PRESSURE**

**COLD & FLU**

Acetaminophen, Doxylamine Succinate,  
Dextromethorphan HBr

Headache, Fever,  
Sore Throat,  
Minor Aches & Pains

Sneezing, Runny Nose

Cough

**SUGAR &  
ALCOHOL  
FREE**

DECONGESTANT FREE

6 FL OZ (177 mL)

90264154





90407196

VICKS

**TAMPER EVIDENT:**  
Do not use if printed shrinkband seal  
around the neck is broken or missing  
← PEEL BACK FOR DRUG FACTS

**Drug Facts**

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine

**PARENTS:**

Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

DIST. BY: PROCTER & GAMBLE,  
CINCINNATI, OH 45202  
Patents: [www.pg.com/patents](http://www.pg.com/patents)  
[vicks.com](http://vicks.com)

**P&G**  
[www.pg.com](http://www.pg.com)  
90264155



<b>Drug Facts</b>	
<b>Active ingredients (in each 30mL)</b>	<b>Purpose</b>
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
<b>Uses</b> temporarily relieves common cold/flu symptoms:	
<ul style="list-style-type: none"> <li>cough due to minor throat &amp; bronchial irritation</li> <li>sore throat</li> <li>headache</li> <li>minor aches &amp; pains</li> <li>fever</li> </ul>	
<b>Warnings</b>	
<b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take:	
<ul style="list-style-type: none"> <li>more than 4 doses (30 mL each dose) in 24 hrs, which is the maximum daily amount for this product</li> <li>with other drugs containing acetaminophen</li> <li>3 or more alcoholic drinks every day while using this product</li> </ul>	
<b>Allergy Alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include:	
<ul style="list-style-type: none"> <li>skin reddening</li> <li>blisters</li> <li>rash</li> </ul>	
If a skin reaction occurs, stop use and seek medical help right away	
<b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
<b>Do not use</b>	
<ul style="list-style-type: none"> <li>with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> </ul>	
<b>Ask a doctor before use if you have</b>	
<ul style="list-style-type: none"> <li>liver disease</li> <li>cough that occurs with too much phlegm (mucus)</li> <li>persistent or chronic cough such as occurs with smoking, asthma, or emphysema</li> </ul>	
<b>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin</b>	
<b>Stop use and ask a doctor if</b>	
<ul style="list-style-type: none"> <li>pain or cough gets worse or lasts more than 7 days</li> <li>fever gets worse or lasts more than 3 days</li> <li>redness or swelling is present</li> <li>new symptoms occur</li> <li>cough comes back or occurs with rash or headache that lasts.</li> </ul>	
These could be signs of a serious condition.	
<b>If pregnant or breast-feeding,</b> ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
<b>Directions</b>	
<ul style="list-style-type: none"> <li>take only as directed</li> <li>only use the dose cup provided</li> <li>do not exceed 4 doses per 24 hrs</li> </ul>	
adults & children 12 yrs & over	30 mL per dose every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use
<b>Other information</b>	
<ul style="list-style-type: none"> <li>each 30 mL dose contains: sodium 29 mg</li> <li>store at no greater than 25 °C (77 °F) and do not refrigerate</li> </ul>	
<b>Inactive ingredients</b> citric acid, FD&C yellow no. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum	

