

Top Layer

HealthA2Z[®] NDC 69168-473-62
 *Compare to the active ingredient in MiraLAX[®].

Lax Relief

Unflavored Powder/Grit Free

Polyethylene Glycol 3350 Powder for Oral Solution Osmotic Laxative

- Relieves Occasional Constipation/Irregularity
- Softens Stool

30 Once-Daily Doses
 NET WT 17.9 OZ (510 g)

TRUST IN HEALTH
 QUALITY ASSURED
 TRUST IN HEALTH A2Z

Peel here

Top Layer Back

Drug Facts	
Active ingredient (in each dose)	Purpose
Polyethylene Glycol 3350, 17 g (cap filled to line).....	Osmotic Laxative
Use	
<ul style="list-style-type: none"> relieves occasional constipation (irregularity) generally produces a bowel movement in 1 to 3 days 	
Warnings	
Allergy alert: Do not use if you are allergic to polyethylene glycol	
Do not use if you have kidney disease, except under the advice and supervision of a doctor	
Ask a doctor before use if you have	
<ul style="list-style-type: none"> nausea, vomiting or abdominal pain a sudden change in bowel habits that lasts over 2 weeks 	

2104113 1

Second Layer

Drug Facts (continued)
<ul style="list-style-type: none"> irritable bowel syndrome
Ask a doctor or pharmacist before use if you are taking a prescription drug
When using this product you may have loose, watery, more frequent stools
Stop use and ask a doctor if
<ul style="list-style-type: none"> you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. you get diarrhea you need to use a laxative for longer than 1 week
If pregnant or breast-feeding , ask a health professional before use.
Keep out of reach of children.

2

Please see back label for continued Drug Facts

Top Layer Back

Drug Facts (continued)
In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
<ul style="list-style-type: none"> do not take more than directed unless advised by your doctor the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap) adults and children 17 years of age and older: <ul style="list-style-type: none"> use once a day fill to top of white section in cap which is marked to indicate the correct dose (17 g) stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink do not combine with starch-based thickeners used for difficulty swallowing

3

Second Layer

Drug Facts (continued)
<ul style="list-style-type: none"> ensure that the powder is fully dissolved before drinking do not drink if there are any clumps do not use more than 7 days children 16 years of age or under: ask a doctor
Other information
<ul style="list-style-type: none"> store at 20° to 25°C (68° to 77°F) tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken
Inactive ingredients
none
Questions or comments?
Call 1-888-952-0050 (Monday-Friday 9AM-5PM EST)

4

Top Layer

Polyethylene Glycol 3350 Powder for Oral Solution increases frequency of bowel movements and softens the stool.

***This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of MiraLax[®].**

- Dissolves in any Beverage**
- Sugar Free**

Manufactured for: **Allegiant Health** LB2271 R0125
 Deer Park, NY 11729

TS/DRUGS/24/2016 2104112

Unvarnish area for Batch coding details 53 x 35 mm

3 69168 47355 3

Peel here