

4 Chewables



Drug Facts (continued)

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- each tablet contains: calcium 140 mg, magnesium 9 mg
- salicylate 101 mg
- very low sodium
- avoid excessive heat (over 104°F or 40°C)

Directions

- chew or dissolve in mouth
- adults and children 12 years and over: 2 tablets every 1/2 to 1 hour as needed
- do not exceed 8 doses (16 tablets) in 24 hours
- use until diarrhea stops but not more than 2 days

Inactive ingredients

calcium carbonate, D&C Red No. 27 aluminum lake, flavor, magnesium stearate, mannitol, povidone, saccharin sodium, talc

Questions?

1-800-717-3786

Made in Mexico by Procter & Gamble Manufactura S. de R.L. de C.V.
www.pepto-bismol.com

Product repackaged and distributed with permission of manufacturer by:
Lil' Drug Store Products, Inc., 9300 Earhart Lane SW, Cedar Rapids, IA 52404
1-877-507-6516 M-F (8 AM-4:30 PM CST) 97232C-US-02-13-A

Pepto-Bismol™

BISMUTH SUBSALICYLATE
UPSET STOMACH RELIEVER/ANTIDIARRHEAL

5 SYMPTOM Digestive Relief

- heartburn
- indigestion
- nausea
- upset stomach
- diarrhea



4
Chewable
Tablets





TAMPER EVIDENT:
Do not use if the printed inner wrap is broken or missing.

Drug Facts

Active ingredient (in each tablet) **Purposes**

Bismuth subsalicylate 262 mg.....Upset stomach reliever and antidiarrheal

Uses relieves

- travelers' diarrhea • diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn • indigestion • nausea • gas • belching • fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer • a bleeding problem • bloody or black stool

Ask a doctor before use if you have • fever • mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for

- anticoagulation (thinning the blood) • diabetes • gout • arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

LIFT PANEL FOR CONTINUED DRUG FACTS