



**Alka-Seltzer Plus Severe Cold PowerFast Fizz Day & Night  
Effervescent Tablets  
Co-Packaged Product**

**The following statement should appear above the Drug Facts box:  
Do not take these products at the same time.**

**The product name should appear outside the Drug Facts box:**

**Alka-Seltzer Plus® Severe Cold PowerFast Fizz Day Effervescent Tablets**

<p><b>Drug Facts</b></p> <table border="0"> <tr> <td><b>Active ingredients (in each tablet)</b></td> <td><b>Purposes</b></td> </tr> <tr> <td>Aspirin 325 mg (NSAID)*.....</td> <td>Pain reliever/fever reducer</td> </tr> <tr> <td>Dextromethorphan hydrobromide 10 mg.....</td> <td>Cough suppressant</td> </tr> <tr> <td>Phenylephrine bitartrate 7.8 mg.....</td> <td>Nasal decongestant</td> </tr> <tr> <td colspan="2">*nonsteroidal anti-inflammatory drug</td> </tr> </table>	<b>Active ingredients (in each tablet)</b>	<b>Purposes</b>	Aspirin 325 mg (NSAID)*.....	Pain reliever/fever reducer	Dextromethorphan hydrobromide 10 mg.....	Cough suppressant	Phenylephrine bitartrate 7.8 mg.....	Nasal decongestant	*nonsteroidal anti-inflammatory drug		<p><b>Annotations</b></p>
<b>Active ingredients (in each tablet)</b>	<b>Purposes</b>										
Aspirin 325 mg (NSAID)*.....	Pain reliever/fever reducer										
Dextromethorphan hydrobromide 10 mg.....	Cough suppressant										
Phenylephrine bitartrate 7.8 mg.....	Nasal decongestant										
*nonsteroidal anti-inflammatory drug											
<p><b>Uses</b></p> <ul style="list-style-type: none"> <li>temporarily relieves these symptoms due to a cold with cough:             <ul style="list-style-type: none"> <li>minor aches and pains</li> <li>headache</li> <li>cough</li> <li>sinus congestion and pressure</li> <li>nasal congestion</li> <li>sore throat</li> </ul> </li> <li>temporarily reduces fever</li> </ul>											
<p><b>Warnings</b></p> <p><b>Reye's syndrome:</b> Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p><b>Allergy alert:</b> Aspirin may cause a severe allergic reaction which may include:</p> <ul style="list-style-type: none"> <li>hives</li> <li>facial swelling</li> <li>asthma (wheezing)</li> <li>shock</li> </ul> <p><b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you</p> <ul style="list-style-type: none"> <li>are age 60 or older</li> <li>have had stomach ulcers or bleeding problems</li> <li>take a blood thinning (anticoagulant) or steroid drug</li> <li>take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</li> <li>have 3 or more alcoholic drinks every day while using this product</li> <li>take more or for a longer time than directed</li> </ul> <p><b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p>											
<p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>if you are allergic to aspirin or any other pain reliever/fever reducer</li> <li>if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> <li>if you have ever had an allergic reaction to this product or any of its ingredients</li> <li>in children under 12 years of age</li> </ul>											
<p><b>Ask a doctor before use if</b></p> <ul style="list-style-type: none"> <li>stomach bleeding warning applies to you</li> <li>you have a history of stomach problems, such as heartburn</li> <li>you have high blood pressure, heart disease, liver cirrhosis, or kidney disease</li> <li>you are taking a diuretic</li> <li>you have             <ul style="list-style-type: none"> <li>asthma</li> <li>thyroid disease</li> <li>diabetes</li> <li>cough that occurs with excessive phlegm (mucus)</li> <li>difficulty in urination due to enlargement of the prostate gland</li> <li>persistent or chronic cough such as occurs with smoking, asthma, or emphysema</li> <li>a sodium-restricted diet</li> </ul> </li> </ul>											





# Alka-Seltzer Plus Severe Cold PowerFast Fizz Night Effervescent Tablets

## Drug Facts

### Active ingredients (in each tablet)

Aspirin 325 mg (NSAID)\*.....Pain reliever/fever reducer  
 Dextromethorphan hydrobromide 10 mg.....Cough suppressant  
 Doxylamine succinate 6.25 mg.....Antihistamine  
 Phenylephrine bitartrate 7.8 mg.....Nasal decongestant  
 \*nonsteroidal anti-inflammatory drug

### Purposes

### Uses

- temporarily relieves these symptoms due to a cold:
  - minor aches and pains
  - sinus congestion and pressure
  - sore throat
  - temporarily reduces fever
  - headache
  - cough
  - nasal congestion
  - runny nose
  - sneezing

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use to sedate children.**

### Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have
  - asthma
  - diabetes
  - thyroid disease
  - glaucoma
  - cough that occurs with excessive phlegm (mucus)
  - a breathing problem such as emphysema or chronic bronchitis
  - difficulty in urination due to enlargement of the prostate gland
  - persistent or chronic cough such as occurs with smoking, asthma, or emphysema
  - a sodium-restricted diet

