

Drug Facts (continued)

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 14 mg
- store at no greater than 25°C

Inactive ingredients citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions? 1-800-362-1683

Drug Facts (continued)

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema

When using this product, do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
 - symptoms do not improve within 7 days or occur with a fever
 - cough persists for more than 7 days, comes back or occurs with a fever, rash or persistent headache
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.



Temper Evident:
Do not use if printed shrinkband seal
around the neck is broken or missing
← PEEL BACK FOR DRUG FACTS

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 100 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves common cold symptoms:
 - nasal congestion
 - sinus congestion & pressure
 - cough due to minor throat & bronchial irritation
 - cough to help you sleep
 - reduces swelling of nasal passages
 - temporarily restores freer breathing through the nose
 - promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

DIST. BY: PROCTER & GAMMA
CINCINNATI, OH 45202
Patents: www.pg.com/
www.vicks.com

P&G
www.pg.com
91767258

PLASTIC BOTTLE

23900 04269 8

3

VICKS®

DayQuil Kids™

HONEY

**COLD & COUGH
+ MUCUS**

Guaifenesin, Phenylephrine HCl,
Dextromethorphan HBr

Chest Congestion
Nasal Congestion
Cough

AGES
6+

Drug Facts (continued)

- cough persists for more than 7 days, comes back or occurs with a fever, rash, or persistent headache
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	do not use unless directed by a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 14 mg
- store at no greater than 25°C

Inactive ingredients citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions? 1-800-362-1683

Drug Facts (continued)

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- **do not use more than directed.**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not improve within 7 days or occur with a fever



Tamper Evident:
 Do not use if printed shrinkband seal
 around the neck is broken or missing
 ← PEEL BACK FOR DRUG FACTS

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Dextromethorphan HBr 10 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine
Phenylephrine HCl 5 mg	Nasal decongestant

Uses temporarily relieves common cold symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

DIST. BY: PROCTER & GAMBLE,
 CINCINNATI, OH 45202
 Patents: www.pg.com/patents
www.vicks.com

P&G
www.pg.com
 91786873

