


Bayer Cross Current Size: 0.308"
Cross Meets Ratio Requirements of 7mm (.2755") or larger
(5mm for blisters/small pouches)
1/4th of Cross required Clear Space (.308") = 0.077"



ARTWORK

SGK
 18211 NE 68th Street,
 E120 Redmond, WA 98052
 T: 425-881-5454
 SGK is a Matthews International Corporation

Job #: 402720215 **Version:** r1
Date: 10FEB2022
Printer: WestRock
Printer Location: East Flat Rock, NC

Client: Bayer HealthCare/USA Consumer Healthcare
Project Type: Production Art

Brand: Coricidin
Bayer#/SBrnd/Dosage: CORICIDIN HBP Cold & Flu Tablet 24 Each Carton PCC0020321

Item#: 88764994 **Die:** BY2144
EPC (EAN/UPC): 041100597885 **UPC Specs:** M: 100% H: 0.84'
C12B/12of5/2DMatrix/Pharma: 4994 **Software:** Illustrator 2020
File Name: 88764994_CORICIDIN_HBP_ColdFlu_Tab_24ea_ctn_r1

Printing Inks

1	PMS Cool Gray 11 C	5	Magenta
2	PMS 186 C	6	Yellow
3	PMS 7579 C		White 1
4	Black		White 2
			Varnish

Non-Printing

- Dieline
- Substrate

Method of Printing:

- combo
- flexo
- offset
- screen
- gravure
- hot stamp
- digital
- dry offset

DRUG FACTS INFO <small>Updated 5/18/18</small>		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt.**	9 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold/italic*	8 pt.
Subheadings	>6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	5 pt.
Leading	>0.5 pt.	0.5 pt.
Hairlines	0.5 pt. rule	0.5 pt.
Box and Barlines†	>0.5 pt. rule	1 pt.
Characters Per Inch:	No more than 39	<39

* Headings are to be at least 2pts larger than Body Text.
 ** Drug Facts Title to be larger than all other type size used in DFP
 † FDA Recommends 2-Spot rule

All Images/Codes are FINAL NOTE: THIS COLOR PROOF INDICATES COLOR BREAK ONLY AND MAY NOT ACCURATELY REFLECT ACTUAL PRODUCTION COLOR.



Coricidin® HBP

Night formula

Drug Facts

Active ingredients (in each capsule)	Purposes
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan hydrobromide 10 mg.....	Cough suppressant
Doxylamine succinate 6.25 mg.....	Antihistamine

Uses

- temporarily relieves these symptoms due to a cold or flu:
 - minor aches and pains
 - headache
 - cough
 - sore throat
 - runny nose
 - sneezing
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age



Coricidin[®] HBP

Ask a doctor before use if you have

- liver disease • glaucoma
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- may cause marked drowsiness
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts.
- These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- do not take the Day and Night products at the same time; wait 4 hours after taking the last Day dose before taking the Night product.
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 4 capsules in 12 hours or as directed by a doctor.
- children under 12 years of age: do not use

Other information

- store at 15°-25°C (59°-77°F)

Inactive ingredients D&C yellow No.10, FD&C blue No.1, gelatin, glycerin, lecithin, medium-chain triglycerides, polyethylene glycol, potassium aluminum silicate, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, soybean oil, titanium dioxide

Questions or comments? 1-800-317-2165 (Mon-Fri 9AM -5PM EST)