



Cetirizine

Hydrochloride Tablets USP, 10 mg
Antihistamine

ALLERGY

Dr.Reddy's

NDC 55111-699-30

Compare to the active ingredient in Zyrtec® Tablets*

1 Bottle Inside

Original Prescription Strength

Cetirizine

Hydrochloride Tablets USP, 10 mg
Antihistamine

ALLERGY

24
HOUR

- Relief of:
- Sneezing
 - Runny Nose
 - Itchy, Watery Eyes
 - Itchy Throat or Nose

Indoor &
Outdoor
Allergies



Actual Size

30 Tablets

TAMPER EVIDENT: DO NOT USE IF
FOIL SEAL UNDER CAP PRINTED
WITH "SEALED FOR YOUR
PROTECTION" IS BROKEN OR MISSING

*This product is not manufactured or
distributed by McNeil Consumer Healthcare,
distributor of Zyrtec® Tablets. Zyrtec® is a
registered trademark of Johnson & Johnson.

DISTRIBUTED BY:
Dr. Reddy's Laboratories, Inc.
Princeton, NJ 08540

Made in India Rev: 04/2024

Drug Facts

| Active ingredient (in each tablet) | Purpose |
|------------------------------------|---------------|
| Cetirizine HCl USP, 10 mg | Antihistamine |

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.
Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product ■ drowsiness may occur ■ avoid alcoholic drinks
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness
 ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.
If pregnant or breast-feeding:
 ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

| | |
|--|--|
| adults and children 6 years and over | one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms. |
| adults 65 years and over | ask a doctor |
| children under 6 years of age | ask a doctor |
| consumers with liver or kidney disease | ask a doctor |

Other information
 ■ store between 20° to 25°C (68° to 77°F)

Inactive ingredients
 hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, povidone, starch, titanium dioxide.

Questions? Call 1-888-375-3784



LOT

EXP

| JOB INFORMATION | HISTORY | COLORS | Structure Reference |
|--------------------------------------|-------------------|---------|---------------------|
| ITEM#: 150100559_CL_CET10mg_30ct_IFC | HR Rev-0: 4-10-24 | Rev-6: | Structure Reference |
| DIELINE#: 1055 | HR Rev-1: 4-25-24 | Rev-7: | Varnish |
| | Rev-2: | Rev-8: | Measurement info. |
| | Rev-3: | Rev-9: | |
| | Rev-4: | Rev-10: | |
| | Rev-5: | Rev-11: | |