

BLACK  
ELDERBERRY  
FLAVORED

VICKS



DayQuil™

ELDERBERRY  
COLD & FLU

Acetaminophen,  
Dextromethorphan HBr



Headache,  
Fever, Sore Throat,  
Minor Aches & Pains  
Cough

12 FL OZ (354 mL)

Non-Drowsy



21022682

BLACK  
RIBBERRY  
FLAVORED

21034428

BLACK  
ELDERBERRY  
FLAVORED

VICKS

TAMPER EVIDENT:  
DO NOT USE IF PRINTED SHRINKBAND  
ON BOTTLE IS BROKEN OR MISSING.  
← PEEL BACK FOR DRUG FACTS

**Drug Facts**

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant

**Uses** temporarily relieves common cold/flu symptoms:  
• cough due to minor throat & bronchial irritation  
• sore throat • headache • minor aches & pains • fever

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if  
• adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product  
• child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product  
• taken with other drugs containing acetaminophen  
• adult has 3 or more alcoholic drinks every day while using this product  
**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash  
If a skin reaction occurs, stop use and seek medical help right away

DIST. BY: PROCTER & GAMBLE,  
CINCINNATI, OH 45202  
Patents: [www.pg.com/patents](http://www.pg.com/patents)

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**P&G**  
21007712

Empty & Recycle Caps

PLASTIC BOTTLE PLASTIC CUP

3 23900 03340 5

## Drug Facts

### Active ingredients (in each 15 mL) Purpose

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg.....Cough suppressant

**Uses** temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat • headache • minor aches & pains • fever

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
- child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include: • skin reddening • blisters • rash  
If a skin reaction occurs, stop use and seek medical help right away

- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

### Other information

- each 15 mL contains: sodium 13 mg
- store at no greater than 25°C and do not refrigerate

**Inactive ingredients** citric acid, FD&C Blue No. 1, FD&C Red No. 40, flavor (with elderberry), glycerin, propylene glycol, saccharin sodium, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, water, xanthan gum

**Questions?** 1-800-362-1683

## Drug Facts (continued)

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- liver disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin.

### Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.