


McNEIL P/N 8737741
CTN TY ARTH PAIN CPLT 24 P&T



EXP: 3 0045-0838-247



See New Warnings Information

NDC 50580-112-24

Push & Turn Cap

TYLENOL
ARTHRITIS PAIN
Acetaminophen extended-release tablets *Pain Reliever/Fever Reducer*
For The Temporary Relief Of Minor Arthritis Pain

24 CAPLETS 650 mg each

Caplets[®]
*Capsule-Shaped, Bi-Layer Tablets

TYLENOL
ARTHRITIS PAIN
Acetaminophen extended-release tablets Pain Reliever/Fever Reducer
For The Temporary Relief Of Minor Arthritis Pain

24 CAPLETS 650 mg each *Capsule-Shaped, Bi-Layer Tablets

Caplets[®]

Contains No Aspirin

Drug Facts (continued)

Other information

■ store at 20-25°C (68-77°F). Avoid excessive heat 40°C (104°F).

■ do not use if carton is opened or neck wrap or foil inner seal imprinted with "Safety Seal®" is broken

■ see end panel for lot number and expiration date

Inactive ingredients

camellia wax, corn starch, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin


Questions or comments?

call 1-877-895-3665

Manufactured by:
McNeil Healthcare LLC
LAS PIEDRAS, PUERTO RICO 00771
For:
McNeil Consumer Healthcare
DIVISION OF MCNEIL-PPC, INC.
FORT WASHINGTON, PA 19034 USA
© MCN-PPC, INC. 2010
Visit us at www.tylenol.com or call toll-free
1-877-TYLENOL (1-877-895-3665)



08737741



YELLOW ORANGE BLUE RED BLK

APPROVED FOR RELEASE TO PRODUCTION

by: **GREAT ATLANTIC** as authorized by: **McNeil**

Approver #1

Signature:

Name: Meredith Faragalli

Title: Production Coordinator

Date: 7/19/2010

Approver #2

Signature:

Name: Diane O'Dwyer

Title: Packaging Graphics Manager

Date: 7/19/2010

Copy & content compared to original document no. LCC 2010-0460

08737741

Drug Facts (continued)

Stop use and ask a doctor if

■ pain gets worse or lasts more than 10 days

■ fever gets worse or lasts more than 3 days

■ new symptoms occur

■ redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

(See overdose warning)

■ do not take more than directed

adults

■ take 2 caplets every 8 hours with water

■ swallow whole – do not crush, chew, split or dissolve

■ do not take more than 6 caplets in 24 hours

■ do not use for more than 10 days unless directed by a doctor

■ ask a doctor

under 18 years of age

8737741