

20 TABLETS

# XL-3\*

## COLD MEDICINE

Fast and strong relief of:

- NASAL CONGESTION • RUNNY NOSE
- FEVER • ACHES & PAINS • SNEEZING

SEE NEW WARNINGS INFORMATION

### Drug Facts

#### Active ingredient (in each tablet)

	Purposes
Acetaminophen 325mg.....	Pain reliever/fever reducer
Chlorpheniramine maleate 2mg.....	Antihistamine
Phenylephrine HCl 5 mg.....	Nasal decongestant

#### Uses

temporarily relieves

- minor aches and pains ■ headaches ■ nasal congestion ■ sinus congestion & pressure ■ runny nose ■ sneezing

#### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 tablets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen

- 3 or more alcoholic drinks every day while using this product

#### Do not use

- for more than 7 days
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have** ■ liver disease ■ heart disease ■ glaucoma ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis

**Ask a doctor or pharmacist before use if you** ■ are taking the blood thinning drug warfarin ■ are taking sedatives or tranquilizers

**When using this product do not use more than directed** ■ excitability may occur, especially in children ■ may cause drowsiness ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if** ■ redness or swelling is present ■ new symptoms occur ■ you get nervous, dizzy, or sleepless ■ pain or nasal congestion gets worse or lasts for more than 7 days ■ fever gets worse or lasts for more than 3 days

- You may report side effects to 1-888-952-0050

**If pregnant or breast-feeding,** ask a health professional before use

**Keep out of reach of children.** In case of overdose, contact a Poison Control Center immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms

### Drug Facts (continued)

**Directions** ■ do not use more than directed ■ Adults and children 12 years of age and older: take 2 tablets every 4 hours ■ Do not take more than 12 tablets in 24 hours ■ Children under 12 years of age: ask a doctor

**Other information** ■ store at room temperature in a dry place ■ do not use if blister package is torn

**Inactive ingredients** corn starch, hydroxypropyl methylcellulose, microcrystalline cellulose, polyethylene glycol, stearic acid



Distributed by: Seldar, SA de CV  
1807 Britton, Houston, TX 77043  
\*Trademark  
LB1108  
R0512