

Drug Facts (continued)

• store at controlled room temperature 20-25°C (68-77°F). Protect from moisture.

Inactive ingredients: acacia, alginate acid, carnauba wax, croscoll silicon dioxide, dibasic calcium phosphate, FD&C blue no. 1 aluminum lake, iron oxide black, magnesium stearate, microcrystalline cellulose, polybutyl methacrylate, polyethylene glycol, polyethylene glycol stearate, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, talc, titanium dioxide

Questions? call 1-800-482-0061

**GENTLE OVERNIGHT
RELIEF YOU CAN TRUST**

24 PILLS



STIMULANT LAXATIVE

SENNOSIDES, USP, 25 mg

exlax

NDC 0067-0016-24

MAXIMUM STRENGTH

exlax

24 PILLS

MAXIMUM STRENGTH

MAXIMUM STRENGTH

exlax

EXP.

LOT.

Tamper Evident Feature: Ex-Lax® Pills
are sealed in blister packets. Use only
if the individual seal is unbroken.

The Ex-Lax® Guarantee: When taken as directed, Ex-Lax® is guaranteed to work gently and effectively overnight or your money back. Return product to Novartis, Novartis Consumer Health, Inc. 2007675

©2012

Novartis Consumer Health, Inc.
Persippany, NJ 07054-0822

3 0067-0016-24 1



Drug Facts

Active ingredient (in each pill)

Sennosides, USP, 25 mg

Purpose

Stimulant laxative

Uses • relieves occasional constipation (irregularity) • generally produces bowel movement in 6 to 12 hours

Warnings

Do not use laxative products when abdominal pain, nausea or vomiting are present unless directed by a doctor

Ask a doctor before use if you have noticed a sudden change in bowel habits that persists over a period of 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Laxatives may affect how other drugs work. Take this product 2 or more hours before or after other drugs.

When using this product do not use for a period longer than 1 week.

Stop use and ask a doctor if rectal bleeding or failure to have a bowel movement occur after use of a laxative. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

• swallow pill(s) with a glass of water • swallow pill(s) whole, do not crush, break or chew

adults and children 12 years of age and older 2 pills once or twice daily

children 6 to under 12 years of age 1 pill once or twice daily

children under 6 years of age ask a doctor

Other information

• each pill contains: calcium 50 mg • sodium free