

**Drug Facts****Active ingredient (in each tablet)**

Loratadine 5 mg

**Purpose**

Antihistamine

**Uses**

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

**Warnings****Do not use** if you have ever had an allergic reaction to this product or any of its ingredients.**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.**When using this product** do not take more than directed. Taking more than directed may cause drowsiness.**Stop use and ask a doctor** if an allergic reaction to this product occurs. Seek medical help right away.**If pregnant or breast-feeding**, ask a health professional before use.**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.**Directions**

- place 1 tablet on tongue; tablet disintegrates, with or without water
- |  |  |
|--|--|
| adults and children 6 years and over   | 1 tablet every 12 hours; not more than 2 tablets in 24 hours |
| children under 6 years of age          | ask a doctor   |
| consumers with liver or kidney disease | ask a doctor   |

**Drug Facts (continued)****Other information**

- safety sealed: do not use if the individual blister unit imprinted with Claritin® RediTabs® is open or torn
- store between 20° to 25° C (68° to 77° F)
- use tablet immediately after opening individual blister
- complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration

**Inactive ingredients**

anhydrous citric acid, gelatin, mannitol, mint flavor

**Questions or comments?**1-888-CLARITIN (1-800-252-7434) or [www.claritin.com](http://www.claritin.com)**Claritin®**  
**RediTabs®****12**  
**Hour**

• No Water Needed • Melts in your mouth

Follow these directions carefully. Do not attempt to push the tablet through the foil.



1. Peel back outer edge.



2. Gently push tablet out.



3. Place the tablet on tongue and close mouth. The tablet will disintegrate.