



<p>Drug Facts</p> <p>Active ingredient (in each chewable tablet) Purpose</p> <p>Cetirizine HCl 2.5 mg Antihistamine</p>	<p>Drug Facts (continued)</p> <p>Directions</p> <p>■ may be taken with or without water</p> <p>■ chew or crush tablets completely before swallowing</p>								
<p>Uses</p> <p>temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:</p> <ul style="list-style-type: none"> ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat 	<table border="1"> <tr> <td>children 2 to under 6 years of age</td> <td>Chew and swallow 1 tablet (2.5 mg) once daily; if needed, dose can be increased to a maximum of 2 tablets (5 mg) once daily or 1 tablet (2.5 mg) every 12 hours. Do not give more than 2 tablets (5 mg) in 24 hours.</td> </tr> </table>	children 2 to under 6 years of age	Chew and swallow 1 tablet (2.5 mg) once daily; if needed, dose can be increased to a maximum of 2 tablets (5 mg) once daily or 1 tablet (2.5 mg) every 12 hours. Do not give more than 2 tablets (5 mg) in 24 hours.						
children 2 to under 6 years of age	Chew and swallow 1 tablet (2.5 mg) once daily; if needed, dose can be increased to a maximum of 2 tablets (5 mg) once daily or 1 tablet (2.5 mg) every 12 hours. Do not give more than 2 tablets (5 mg) in 24 hours.								
<p>Warnings</p> <p>Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.</p> <p>Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.</p> <p>Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.</p>	<table border="1"> <tr> <td>adults and children 6 years and over</td> <td>Chew and swallow 2 tablets (5 mg) or 4 tablets (10 mg) once daily depending upon severity of symptoms; do not take more than 4 tablets (10 mg) in 24 hours.</td> </tr> <tr> <td>adults 65 years and over</td> <td>Chew and swallow 2 tablets (5 mg) once daily; do not take more than 2 tablets (5 mg) in 24 hours.</td> </tr> <tr> <td>children under 2 years of age</td> <td>ask a doctor</td> </tr> <tr> <td>consumers with liver or kidney disease</td> <td>ask a doctor</td> </tr> </table>	adults and children 6 years and over	Chew and swallow 2 tablets (5 mg) or 4 tablets (10 mg) once daily depending upon severity of symptoms; do not take more than 4 tablets (10 mg) in 24 hours.	adults 65 years and over	Chew and swallow 2 tablets (5 mg) once daily; do not take more than 2 tablets (5 mg) in 24 hours.	children under 2 years of age	ask a doctor	consumers with liver or kidney disease	ask a doctor
adults and children 6 years and over	Chew and swallow 2 tablets (5 mg) or 4 tablets (10 mg) once daily depending upon severity of symptoms; do not take more than 4 tablets (10 mg) in 24 hours.								
adults 65 years and over	Chew and swallow 2 tablets (5 mg) once daily; do not take more than 2 tablets (5 mg) in 24 hours.								
children under 2 years of age	ask a doctor								
consumers with liver or kidney disease	ask a doctor								
<p>When using this product</p> <ul style="list-style-type: none"> ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery 	<p>Other information</p> <ul style="list-style-type: none"> ■ store between 20° to 25°C (68° to 77°F) ■ do not use if blister unit is torn or broken 								
<p>Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.</p> <p>If pregnant or breast-feeding:</p> <ul style="list-style-type: none"> ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use. 	<p>Inactive ingredients betadex, corn starch, flavor, lactose monohydrate, magnesium stearate, mannitol, silicified microcrystalline cellulose, sucralose</p>								
<p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) →</p>	<p>Questions? call 1-800-343-7805 (toll-free) or 215-273-8755 (collect)</p>								