

ADHERE TO SHOWN BLEED AREAS, TO AVOID ANY BARE SPOTS SHOWING AFTER THE FLAPS ARE GLUED,  
 FOR PHARMACODE AND AREAS WHERE BLACK TEXT IS INDICATED:  
 IF BLACK IS NOT IN COLOUR BREAK, USE DARKEST COLOUR AVAILABLE

BLEED LINE

Compare to the active ingredient in Lamisil AT®\*

Cures Most Athlete's Foot

# Terbinafine Hydrochloride Cream 1% Antifungal Cream

• Relieves Itching & Burning  
• Full Prescription Strength

MPK-6889-0  
0212-0  
M000

T175

NO COPY ON THIS FLAP: LOT & EXPIRY AREA

NO VARNISH ON THIS FLAP

NO VARNISH ON THIS FLAP

NO VARNISH ON THIS FLAP

P/C

Questions? Call 1-866-923-4914

**Drug Facts (continued)**

**Other information** • do not use if seal on tube is broken or is not visible • see carton or tube crimp for lot number and expiration date • store at controlled room temperature 20°-25°C (68°-77°F)

**Inactive ingredients:** benzyl alcohol, cetyl alcohol, cetyl palmitate, isopropyl myristate, polysorbate 60, purified water, sodium hydroxide, sorbitan monostearate, stearyl alcohol.

**Drug Facts**

**Active ingredient** Terbinafine hydrochloride 1% Antifungal Cream 1%

**Uses**

- cures most athlete's foot (tinea pedis)
- cures most jock itch (tinea cruris) and ringworm (tinea corporis)
- relieves itching, burning, cracking and scaling which accompany these conditions

**Warnings**

**For external use only**

**Do not use** • on nails or scalp • in or near the mouth or the eyes • for vaginal yeast infections

**When using this product** do not get into the eyes. If eye contact occurs, rinse thoroughly with water.

**Stop use and ask a doctor if** • too much irritation occurs or gets worse.

**Keep out of reach of children.** If swallowed, get medical help or contact a poison control center right away.

**Directions**

- adults and children 12 years and older
- use the tip of the cap to break the seal and open the tube
- wash the affected skin with soap and water and dry completely before applying
- athlete's foot wear well-fitting, ventilated shoes. Change shoes and socks at least once daily.
- between the toes only: apply twice a day (morning and night) for 1 week or as directed by a doctor.
- on the bottom or sides of the foot: apply twice a day (morning and night) for 2 weeks or as directed by a doctor.
- for jock itch and ringworm: apply once a day (morning or night) for 1 week or as directed by a doctor.
- wash hands after each use
- children under 12 years: ask a doctor

**2 weeks on the bottom or sides of the foot.**

**1 week between the toes.**

Distributed by:

Made in Canada.

This product is not manufactured or distributed by Novartis Consumer Health, Inc., owner of the registered trademark Lamisil AT®.

Antifungal Cream

Terbinafine Hydrochloride Cream 1% Antifungal Cream

NET WT 1 oz (30 g)

Terbinafine Hydrochloride Cream 1% Antifungal Cream

**ATTACHED LEGEND TO ALL DIGITAL MECHANICALS**

**STANDARD FORMAT:**  
 Helvetica Neue 55 Roman  
 100% Horizontal Scale

**Drug Facts - 14pt Helvetica Neue 76 Italic**

**Headings - 8pt Helvetica Neue 76 Italic**

**Sub Heads - 6pt Helvetica Neue 75 Bold**

Text - 6pt Helvetica Neue 55 Roman

**Round Bullets - 6pt Helvetica Neue 55 Roman**

**Border - 2.5 pt Rule, 0.025" gap all around**

**Barline - 2.5 pt Rule** \_\_\_\_\_

**Hairline - .5 pt Rule** \_\_\_\_\_

Certified by: **MS**

PMS BLACK    PMS CYAN    PMS 280    PMS 877

APPROVAL				
DATE	02/3/12	MPK#	6889-0	ART COPY CTRL.#
PASS#	TerbinafineAF30gCarton6889-0	UPC#		PHARMACOD# M000
DEPARTMENT	SIGNATURE	NO	YES	DATE
PROOFING				
ART			APPROVED	By Kevin Weight at 11:35 am, Feb 05, 2012
MARKETING				
ENGINEERING				
Q.C.				
REG.		APPROVED	APPROVED	By Czarina Ochoa at 11:20 am, Feb 07, 2012    By Nancy L. Westcott at 12:26 pm, Feb 07, 2012
MEDICAL OFFICER				
CLIENT				

THIS COLOR PROOF INDICATES COLOR BREAK ONLY AND MAY NOT ACCURATELY REFLECT ACTUAL PRODUCTION COLOR IF YOU HAVE ANY QUESTIONS OR COMMENTS PLEASE CALL TARO PHARMACEUTICALS-ART DEPARTMENT 800-544-1449.